BPS Dendroscopy on Local Calabi-Yau Threefolds

Boris Pioline





< D > < P > < E >

String-Math 2022, 14/07/2022

Work in progress with Pierrick Bousseau, Pierre Descombes and Bruno Le Floch







Dendrochronology

B. Pioline (LPTHE, Paris)

BPS Dendroscopy

▲ ■ ▶ ■ つへの 14/07/2022 2/35

δενδροσκοπια= analyzing the BPS spectrum in terms of attractor flow trees



Attractor flow trees on $K_{\mathbb{P}^2}$, $\gamma = [1, 0, -3)$, $\mathcal{M} = \text{Hilb}_4 \mathbb{P}^2$

• In type IIA string theory compactified on a Calabi-Yau threefold X, the BPS spectrum consists of bound states of D6-D4-D2-D0 branes, described mathematically by objects *E* in the derived category of coherent sheaves $C = D^b Coh(X)$ [Douglas'01]

- In type IIA string theory compactified on a Calabi-Yau threefold X, the BPS spectrum consists of bound states of D6-D4-D2-D0 branes, described mathematically by objects *E* in the derived category of coherent sheaves $C = D^b Coh(X)$ [Douglas'01]
- The BPS index or Donaldson-Thomas invariant Ω_z(γ) counts stable states with charge γ = ch E ∈ H_{even}(X, Q) saturating the BPS bound M(γ) ≥ |Z(γ)|, where Z ∈ Hom(Γ, C) depends on the complexified Kähler moduli z ∈ M.

- In type IIA string theory compactified on a Calabi-Yau threefold X, the BPS spectrum consists of bound states of D6-D4-D2-D0 branes, described mathematically by objects E in the derived category of coherent sheaves $C = D^b Coh(X)$ [Douglas'01]
- The BPS index or Donaldson-Thomas invariant Ω_z(γ) counts stable states with charge γ = ch E ∈ H_{even}(X, Q) saturating the BPS bound M(γ) ≥ |Z(γ)|, where Z ∈ Hom(Γ, C) depends on the complexified Kähler moduli z ∈ M.
- $\Omega_z(\gamma)$ is locally constant on \mathcal{M} , but can jump across real codimension one walls of marginal stability $\mathcal{W}(\gamma_1, \gamma_2) \subset \mathcal{M}$, where the phases of the central charges $Z(\gamma_1)$ and $Z(\gamma_2)$ with $\gamma = m_1\gamma_1 + m_2\gamma_2$ become aligned [Kontsevich Soibelman'08, Joyce Song'08]

< ロ > < 同 > < 回 > < 回 > < 回 > <

- In type IIA string theory compactified on a Calabi-Yau threefold X, the BPS spectrum consists of bound states of D6-D4-D2-D0 branes, described mathematically by objects *E* in the derived category of coherent sheaves $C = D^b Coh(X)$ [Douglas'01]
- The BPS index or Donaldson-Thomas invariant Ω_z(γ) counts stable states with charge γ = ch E ∈ H_{even}(X, Q) saturating the BPS bound M(γ) ≥ |Z(γ)|, where Z ∈ Hom(Γ, C) depends on the complexified Kähler moduli z ∈ M.
- $\Omega_z(\gamma)$ is locally constant on \mathcal{M} , but can jump across real codimension one walls of marginal stability $\mathcal{W}(\gamma_1, \gamma_2) \subset \mathcal{M}$, where the phases of the central charges $Z(\gamma_1)$ and $Z(\gamma_2)$ with $\gamma = m_1\gamma_1 + m_2\gamma_2$ become aligned [Kontsevich Soibelman'08, Joyce Song'08]
- Physically, multi-centered black hole solutions (dis)appear across the wall [Denef Moore '07, ..., Manschot BP Sen '11].

< ロ > < 同 > < 回 > < 回 > < 回 > <

BPS spectrum on local surfaces

• For a non-compact CY3 of the form $X = K_S$ where S is a complex Fano surface, there is an injection $\iota_* : D^b \operatorname{Coh}(S) \to D^b_c(X)$ lifting an object E with Chern character $\gamma = [r, d, \operatorname{ch}_2]$ to a bound state of r D4-branes wrapped on S, c D2-branes and ch₂ D0-branes.

BPS spectrum on local surfaces

- For a non-compact CY3 of the form $X = K_S$ where *S* is a complex Fano surface, there is an injection $\iota_* : D^b \operatorname{Coh}(S) \to D^b_c(X)$ lifting an object *E* with Chern character $\gamma = [r, d, \operatorname{ch}_2]$ to a bound state of *r* D4-branes wrapped on *S*, *c* D2-branes and ch₂ D0-branes.
- At large volume, the central charge is quadratic in complexified Kähler moduli $z^a = b^a + it^a$,

$$Z(\gamma) \sim -\int_{\mathcal{S}} e^{-z^a H_a} \operatorname{ch} E = -r \, z^a Q_{ab} z^b + z^a d_a - \operatorname{ch}_2$$

 $\Omega_z(\gamma)$ reduces to the Gieseker index $\Omega_\infty(\gamma)$, given (up to sign) by the Euler number of the moduli space of Gieseker semi-stable sheaves on *S* with Chern character γ .

BPS spectrum on local surfaces

- For a non-compact CY3 of the form $X = K_S$ where *S* is a complex Fano surface, there is an injection $\iota_* : D^b \operatorname{Coh}(S) \to D^b_c(X)$ lifting an object *E* with Chern character $\gamma = [r, d, \operatorname{ch}_2]$ to a bound state of *r* D4-branes wrapped on *S*, *c* D2-branes and ch₂ D0-branes.
- At large volume, the central charge is quadratic in complexified Kähler moduli $z^a = b^a + it^a$,

$$Z(\gamma) \sim -\int_{\mathcal{S}} e^{-z^a H_a} \operatorname{ch} E = -r \, z^a Q_{ab} z^b + z^a d_a - \operatorname{ch}_2$$

 $\Omega_z(\gamma)$ reduces to the Gieseker index $\Omega_\infty(\gamma)$, given (up to sign) by the Euler number of the moduli space of Gieseker semi-stable sheaves on *S* with Chern character γ .

 At finite volume, Z receives worldsheet instanton corrections computable by mirror symmetry. Can we determine Ω_Z(γ) anywhere, and understand what are BPS states really "made of" ?

- 2 Kähler moduli space of $K_{\mathbb{P}^2}$
- 3 Attractor flow trees and scattering diagrams
- 4 Large volume scattering diagram
- 5 Towards the exact scattering diagram

- 2 Kähler moduli space of $K_{\mathbb{P}^2}$
- 3 Attractor flow trees and scattering diagrams
- 4 Large volume scattering diagram
- 5 Towards the exact scattering diagram

Kähler moduli space of $K_{\mathbb{P}^2}$

The Kähler moduli space of K_{P²} is the modular curve X₁(3) = ℍ/Γ₁(3) parametrizing elliptic curves with level structure. It admits two cusps LV, C and one elliptic point o of order 3.

Kähler moduli space of $K_{\mathbb{P}^2}$

- The Kähler moduli space of K_{P²} is the modular curve X₁(3) = ℍ/Γ₁(3) parametrizing elliptic curves with level structure. It admits two cusps LV, C and one elliptic point o of order 3.
- The universal cover is parametrized by $au \in \mathbb{H}$:



$$Z_{\tau}(\gamma) = -rT_{D}(\tau) + dT(\tau) - ch_{2}$$
$$T = \int_{\ell} \lambda$$
$$T_{D} = \int_{\ell_{D}} \lambda$$

 λ holomorphic one-form with logarithmic singularities on \mathcal{E}_{τ}

 Since ∂_τλ is holomorphic, its periods are proportional to (1, τ). Integrating on a path from *o* to τ, one finds the Eichler-type integral

$$\begin{pmatrix} T \\ T_D \end{pmatrix} = \begin{pmatrix} 1/2 \\ 1/3 \end{pmatrix} + \int_{\tau_o}^{\tau} \begin{pmatrix} 1 \\ u \end{pmatrix} C(u) \, \mathrm{d}u$$

where $C(\tau) = \frac{\eta(\tau)^9}{\eta(3\tau)^3}$ is a weight 3 modular form for $\Gamma_1(3)$.

 Since ∂_τλ is holomorphic, its periods are proportional to (1, τ). Integrating on a path from *o* to τ, one finds the Eichler-type integral

$$\begin{pmatrix} T \\ T_D \end{pmatrix} = \begin{pmatrix} 1/2 \\ 1/3 \end{pmatrix} + \int_{\tau_0}^{\tau} \begin{pmatrix} 1 \\ u \end{pmatrix} C(u) \, \mathrm{d}u$$

where $C(\tau) = \frac{\eta(\tau)^9}{\eta(3\tau)^3}$ is a weight 3 modular form for $\Gamma_1(3)$.

 This provides an computationally efficient analytic continuation of Z_τ throughout II, and gives access to monodromies:

$$au \mapsto rac{a au + b}{c au + d} = egin{pmatrix} 1 \ T \ T_D \end{pmatrix} \mapsto egin{pmatrix} 1 & 0 & 0 \ m & d & c \ m_D & b & a \end{pmatrix} \cdot egin{pmatrix} 1 \ T \ T_D \end{pmatrix}$$

where (m, m_D) are period integrals of *C* from τ_o to $\frac{a\tau_o - b}{c\tau_o - d}$.

• At large volume, using C = 1 - 9q + ... one finds

$$T = au + \mathcal{O}(q), \quad T_D = \frac{1}{2} au^2 + \frac{1}{8} + \mathcal{O}(q)$$

< D > < P > < E >

• At large volume, using C = 1 - 9q + ... one finds

$$T = \tau + \mathcal{O}(q), \quad T_D = \frac{1}{2}\tau^2 + \frac{1}{8} + \mathcal{O}(q)$$

For τ₂ large enough, one can use the *GL*(2, ℝ)⁺ action on space of Bridgeland stability conditions to absorb the *O*(*q*) corrections:

$$Z_{(s,t)}^{LV}(\gamma) = -\frac{1}{2}(s+it)^2 + d(s+it) - ch_2 ,$$

$$s = \frac{Im T_D}{Im T} , \quad \mu = \frac{d}{r}$$

$$\frac{1}{2}(s^2 + t^2) = -\frac{Im(T\bar{T}_D)}{Im T}$$

$$\mathcal{A} = \{E \stackrel{d}{\rightarrow} F, \mu(E) < s, \mu(F) \ge s\}$$
[Bayer Macri'11]

B. Pioline (LPTHE, Paris)

BPS Dendroscopy

• Near the orbifold point $\tau_o = -\frac{1}{2} + \frac{i}{2\sqrt{3}}$, the BPS spectrum is governed by a quiver with potential:



 The BPS index Ω_τ(γ) coincides with the (signed) Euler number Ω_ζ(γ) of the moduli space of King semi-stable representations of dimension γ = (n₁, n₂, n₃), with FI-parameters θ_i = ImZ_τ(γ_i).

In the chamber θ₁ > 0, θ₃ < 0, the arrows Z_k vanish in any stable representation, and (Q, W) reduces to the Beilinson quiver describing normalized torsion-free sheaves on P²:

$$(n_1) \xrightarrow{X_i} (n_2) \xrightarrow{Y_j} (n_3) \quad \epsilon_{ijk} X^i Y^j = 0$$

Douglas Fiol Romelsberger'00

In the chamber θ₁ > 0, θ₃ < 0, the arrows Z_k vanish in any stable representation, and (Q, W) reduces to the Beilinson quiver describing normalized torsion-free sheaves on P²:

$$(n_1 \xrightarrow{X_i} n_2 \xrightarrow{Y_j} n_3) \quad \epsilon_{ijk} X^i Y^j = 0$$

Douglas Fiol Romelsberger'00

• In [Beaujard BP Manschot'20], we showed that the attractor index $\Omega_{\star}(\gamma) := \Omega_{\langle \gamma, - \rangle}(\gamma)$ vanishes except for $\gamma = \gamma_i$ or $\gamma \propto \gamma_1 + \gamma_2 + \gamma_3$.

In the chamber θ₁ > 0, θ₃ < 0, the arrows Z_k vanish in any stable representation, and (Q, W) reduces to the Beilinson quiver describing normalized torsion-free sheaves on P²:

$$(n_1) \xrightarrow{X_i} (n_2) \xrightarrow{Y_j} (n_3) \quad \epsilon_{ijk} X^i Y^j = 0$$

Douglas Fiol Romelsberger'00

A B > A B >

- In [Beaujard BP Manschot'20], we showed that the attractor index $\Omega_{\star}(\gamma) := \Omega_{\langle \gamma, - \rangle}(\gamma)$ vanishes except for $\gamma = \gamma_i$ or $\gamma \propto \gamma_1 + \gamma_2 + \gamma_3$.
- Moreover, the anti-attractor index Ω_x(γ) := Ω_{-⟨γ,-⟩}(γ) coincide with the Gieseker index Ω_∞(γ), provided −r < d ≤ 0.

In the chamber θ₁ > 0, θ₃ < 0, the arrows Z_k vanish in any stable representation, and (Q, W) reduces to the Beilinson quiver describing normalized torsion-free sheaves on P²:

$$(n_1 \xrightarrow{X_i} n_2 \xrightarrow{Y_j} n_3) \quad \epsilon_{ijk} X^i Y^j = 0$$

Douglas Fiol Romelsberger'00

- In [Beaujard BP Manschot'20], we showed that the attractor index $\Omega_{\star}(\gamma) := \Omega_{\langle \gamma, \rangle}(\gamma)$ vanishes except for $\gamma = \gamma_i$ or $\gamma \propto \gamma_1 + \gamma_2 + \gamma_3$.
- Moreover, the anti-attractor index Ω_x(γ) := Ω_{-⟨γ,-⟩}(γ) coincide with the Gieseker index Ω_∞(γ), provided −r < d ≤ 0.
- A similar conjecture for Ω_{*}(γ) holds for any toric CY3, giving in principle access to DT invariants Ω_ζ(γ) for any ζ ∈ ℝ^{Q0} [Mozgovoy BP'20; Descombes'21]

2) Kähler moduli space of $K_{\mathbb{P}^2}$

3 Attractor flow trees and scattering diagrams

- 4 Large volume scattering diagram
- 5 Towards the exact scattering diagram

The Attractor Flow Tree formula for quivers

The Attractor Flow Tree Formula expresses the BPS index Ω_θ(γ) for any (generic) θ ∈ ℝ^{Q₀} in terms of attractor indices by summing over all possible flow trees: schematically,

$$\Omega_{\theta}(\gamma) \sim \sum_{\gamma = \gamma_1 + \dots + \gamma_n} \left(\sum_{T \in \mathcal{T}_{\theta}(\{\gamma_i\})} \prod_{\nu \in V_T} \langle \gamma_{L(\nu)}, \gamma_{R(\nu)} \rangle \right) \prod_{i=1}^n \Omega_{\star}(\gamma_i)$$

Denef '00; Denef Greene Raugas '01; Denef Moore'07; Manschot '10, Alexandrov BP'18

The Attractor Flow Tree formula for quivers

The Attractor Flow Tree Formula expresses the BPS index Ω_θ(γ) for any (generic) θ ∈ ℝ^{Q0} in terms of attractor indices by summing over all possible flow trees: schematically,

$$\Omega_{\theta}(\gamma) \sim \sum_{\gamma = \gamma_{1} + \dots + \gamma_{n}} \left(\sum_{T \in \mathcal{T}_{\theta}(\{\gamma_{i}\})} \prod_{\nu \in V_{T}} \langle \gamma_{L(\nu)}, \gamma_{R(\nu)} \rangle \right) \prod_{i=1}^{n} \Omega_{\star}(\gamma_{i})$$

Denef '00; Denef Greene Raugas '01; Denef Moore'07; Manschot '10, Alexandrov BP'18

• Here, a flow tree *T* is a binary rooted tree, with edges decorated with charges γ_e , such that $\gamma_v = \gamma_{L(v)} + \gamma_{R(v)}$ at each vertex, with charges γ_i assigned to the leaves and γ to the root.

The Attractor Flow Tree formula for quivers

The Attractor Flow Tree Formula expresses the BPS index Ω_θ(γ) for any (generic) θ ∈ ℝ^{Q₀} in terms of attractor indices by summing over all possible flow trees: schematically,

$$\Omega_{\theta}(\gamma) \sim \sum_{\gamma = \gamma_{1} + \dots + \gamma_{n}} \left(\sum_{T \in \mathcal{T}_{\theta}(\{\gamma_{i}\})} \prod_{\nu \in V_{T}} \langle \gamma_{L(\nu)}, \gamma_{R(\nu)} \rangle \right) \prod_{i=1}^{''} \Omega_{\star}(\gamma_{i})$$

Denef '00; Denef Greene Raugas '01; Denef Moore'07; Manschot '10, Alexandrov BP'18

- Here, a flow tree *T* is a binary rooted tree, with edges decorated with charges γ_e , such that $\gamma_v = \gamma_{L(v)} + \gamma_{R(v)}$ at each vertex, with charges γ_i assigned to the leaves and γ to the root.
- Each edge is embedded in R^{Q₀} along θ_ν = θ_{ρ(ν)} + λ⟨γ_e, -⟩, λ ≥ 0, such that the root vertex maps to θ, and (θ_ν, γ_{L(ν)}) = (θ_ν, γ_{R(ν)}) = 0 at each vertex.

n

Split attractor flows

• The physical picture is that typical multi-centered solutions in $\mathcal{N}=2$ supergravity have a nested structure



Split attractor flows

• The physical picture is that typical multi-centered solutions in $\mathcal{N}=2$ supergravity have a nested structure



• The linear flow in *θ* originates from gradient flow for spherically symmetric black holes [Ferrara Kallosh Strominger'95]

$$r^2rac{\mathrm{d}z^i}{\mathrm{d}r}=-g^{iar{j}}\partial_{ar{j}}|Z(\gamma)|^2$$

Split attractor flows

• The physical picture is that typical multi-centered solutions in $\mathcal{N}=$ 2 supergravity have a nested structure



• The linear flow in *θ* originates from gradient flow for spherically symmetric black holes [Ferrara Kallosh Strominger'95]

$$r^2 rac{\mathrm{d} z^i}{\mathrm{d} r} = -g^{i \overline{j}} \partial_{\overline{j}} |Z(\gamma)|^2$$

At each level ν, the average distance between the clusters of charge γ_L(ν) and γ_R(ν) is fixed, but the orientation in S² gives |(γ_{L(ν)}, γ_{R(ν)})| degrees of freedom. In addition, each center of charge γ_i carries internal degrees of freedom counted by Ω_{*}(γ_i).

• In order to enforce Bose-Fermi statistics whenever two charges coincide, one should replace $\Omega_{\theta}(\gamma)$ by the rational index $\bar{\Omega}_{\theta}(\gamma) = \sum_{d|\gamma} \frac{1}{d^2} \Omega_{\theta}(\frac{\gamma}{d})$ and insert a Boltzmann symmetry factor. [Manschot BP Sen'11]

- In order to enforce Bose-Fermi statistics whenever two charges coincide, one should replace $\Omega_{\theta}(\gamma)$ by the rational index $\bar{\Omega}_{\theta}(\gamma) = \sum_{d|\gamma} \frac{1}{d^2} \Omega_{\theta}(\frac{\gamma}{d})$ and insert a Boltzmann symmetry factor. [Manschot BP Sen'11]
- When the charges γ_i are not linearly independent, some splittings can involve higher valency vertices. One can treat them using the full KS wall-crossing formula, or perturb θ such that only binary trees remain.

- In order to enforce Bose-Fermi statistics whenever two charges coincide, one should replace $\Omega_{\theta}(\gamma)$ by the rational index $\bar{\Omega}_{\theta}(\gamma) = \sum_{d|\gamma} \frac{1}{d^2} \Omega_{\theta}(\frac{\gamma}{d})$ and insert a Boltzmann symmetry factor. [Manschot BP Sen'11]
- When the charges γ_i are not linearly independent, some splittings can involve higher valency vertices. One can treat them using the full KS wall-crossing formula, or perturb θ such that only binary trees remain.
- The attractor flow tree formula is consistent with wall-crossing: the index jumps when *z* crosses the wall $W(\gamma_{L(v_0)}, \gamma_{R(v_0)})$ associated to the primary splitting for one of the trees.

Remarks

• There are additional 'fake walls' where the topology of the trees jump but the total index is constant, thanks to the identity



Remarks

• There are additional 'fake walls' where the topology of the trees jump but the total index is constant, thanks to the identity



• The formula can be refined by replacing

$$\begin{array}{lcl} \langle \gamma_L, \gamma_R \rangle & \to & \frac{y^{\langle \gamma_L, \gamma_R \rangle} - y^{-\langle \gamma_L, \gamma_R \rangle}}{y - 1/y} \\ \bar{\Omega}_{\theta}(\gamma) & \to & \bar{\Omega}_{\theta}(\gamma, y) = \sum_{d \mid \gamma} \frac{y - 1/y}{d(y^d - y^{-d})} \Omega_{\theta}(\frac{\gamma}{d}, y^d) \end{array}$$

Physically, *y* is a fugacity conjugate to angular momentum in \mathbb{R}^3 .

Flow tree formula from scattering diagrams

• For any quiver with potential (Q, W), the scattering diagram \mathcal{D} is the set of real codimension-one rays $\{\mathcal{R}(\gamma), \gamma \in \mathbb{Z}^{Q_0}\}$ defined by [Bridgeland'16]

 $\mathcal{R}(\gamma) = \{\zeta \in \mathbb{R}^{Q_0} : (\zeta, \gamma) = 0, \Omega_{\zeta}(k\gamma) \neq 0 \text{ for some } k \geq 1\}$

Flow tree formula from scattering diagrams

• For any quiver with potential (Q, W), the scattering diagram \mathcal{D} is the set of real codimension-one rays $\{\mathcal{R}(\gamma), \gamma \in \mathbb{Z}^{Q_0}\}$ defined by [Bridgeland'16]

 $\mathcal{R}(\gamma) = \{\zeta \in \mathbb{R}^{Q_0} : (\zeta, \gamma) = 0, \Omega_{\zeta}(k\gamma) \neq 0 \text{ for some } k \geq 1\}$

• Each point along $\mathcal{R}(\gamma)$ is endowed with an automorphism of the quantum torus algebra, (assume γ primitive)

$$\mathcal{U}(\gamma) = \exp(\sum_{m=1}^{\infty} \frac{\bar{\Omega}_{\zeta}(k\gamma, y)}{y^{-1} - y} \mathcal{X}_{k\gamma}) , \quad \mathcal{X}_{\gamma} \mathcal{X}_{\gamma'} = (-y)^{\langle \gamma, \gamma' \rangle} \mathcal{X}_{\gamma + \gamma'}$$



 $\gamma_1 + \gamma_2$ • The WCF ensures that the diagram is consistent, $\prod_{\gamma_i} \mathcal{U}(\gamma_i)^{\pm 1} = 1$ around any codimension 2 intersection. The Attractor Flow Tree Formula determines outgoing rays from incoming rays at each vertex. [Argüz Bousseau '20].

B. Pioline (LPTHE, Paris)

BPS Dendroscopy

Orbifold scattering diagram



B. Pioline (LPTHE, Paris)

э 14/07/2022 19/35

ъ

A 2D slice of the orbifold scattering diagram



B. Pioline (LPTHE, Paris)

14/07/2022 20 / 35

< □ > < □ >

A 2D slice of the orbifold scattering diagram



B. Pioline (LPTHE, Paris)

BPS Dendroscopy

14/07/2022 21 / 35

< A

• More generally, for any $\psi \in \mathbb{R}/2\pi\mathbb{Z}$ define scattering rays as codimension-one loci in the space of Bridgeland stability conditions

 $\mathcal{R}_{\psi}(\gamma) = \{ Z : \operatorname{Re}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) = \boldsymbol{0}, \operatorname{Im}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) > \boldsymbol{0}, \Omega_{\zeta}(\boldsymbol{k}\gamma) \neq \boldsymbol{0} \}$

• More generally, for any $\psi \in \mathbb{R}/2\pi\mathbb{Z}$ define scattering rays as codimension-one loci in the space of Bridgeland stability conditions

 $\mathcal{R}_{\psi}(\gamma) = \{ Z : \operatorname{Re}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) = \boldsymbol{0}, \operatorname{Im}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) > \boldsymbol{0}, \Omega_{\zeta}(\boldsymbol{k}\gamma) \neq \boldsymbol{0} \}$

For a non-compact CY3, Z(γ) is holomorphic in Kähler moduli, thus arg Z(γ) is constant along the gradient flow of |Z(γ)|. Choosing ψ such that z ∈ R_ψ(γ), edges of attractor flow trees lie inside R_ψ(γ_e), while vertices lie in R_ψ(γ_{L(ν)}) ∩ R_ψ(γ_{R(ν)}).

• More generally, for any $\psi \in \mathbb{R}/2\pi\mathbb{Z}$ define scattering rays as codimension-one loci in the space of Bridgeland stability conditions

 $\mathcal{R}_{\psi}(\gamma) = \{ Z : \operatorname{Re}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) = \boldsymbol{0}, \operatorname{Im}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) > \boldsymbol{0}, \Omega_{\zeta}(\boldsymbol{k}\gamma) \neq \boldsymbol{0} \}$

- For a non-compact CY3, Z(γ) is holomorphic in Kähler moduli, thus arg Z(γ) is constant along the gradient flow of |Z(γ)|. Choosing ψ such that z ∈ R_ψ(γ), edges of attractor flow trees lie inside R_ψ(γ_e), while vertices lie in R_ψ(γ_{L(ν)}) ∩ R_ψ(γ_{R(ν)}).
- Besides, since Z(γ) is holomorphic, initial rays must originate from attractor points on the boundary.

- 4 同 ト 4 三 ト 4

• More generally, for any $\psi \in \mathbb{R}/2\pi\mathbb{Z}$ define scattering rays as codimension-one loci in the space of Bridgeland stability conditions

 $\mathcal{R}_{\psi}(\gamma) = \{ Z : \operatorname{Re}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) = \boldsymbol{0}, \operatorname{Im}(\boldsymbol{e}^{-\mathrm{i}\psi}Z(\gamma)) > \boldsymbol{0}, \Omega_{\zeta}(\boldsymbol{k}\gamma) \neq \boldsymbol{0} \}$

- For a non-compact CY3, Z(γ) is holomorphic in Kähler moduli, thus arg Z(γ) is constant along the gradient flow of |Z(γ)|. Choosing ψ such that z ∈ R_ψ(γ), edges of attractor flow trees lie inside R_ψ(γ_e), while vertices lie in R_ψ(γ_{L(ν)}) ∩ R_ψ(γ_{R(ν)}).
- Besides, since Z(γ) is holomorphic, initial rays must originate from attractor points on the boundary.
- Fflow trees are subsets of scattering diagrams, determining sequences of scatterings which produce an outgoing ray R_ψ(γ) passing through the desired point z.

< ロ > < 同 > < 回 > < 回 > < 回 > <

- 2) Kähler moduli space of $K_{\mathbb{P}^2}$
- 3 Attractor flow trees and scattering diagrams
- 4 Large volume scattering diagram
- 5 Towards the exact scattering diagram

• For the large volume stability conditions $Z_{(s,t)}^{LV}$, [Bousseau'19] constructed the scattering diagram \mathcal{D}_{ψ} in (s, t) upper half-plane for $\psi = 0$. For $\psi \neq 0$, just map $(s, t) \mapsto (s - t \tan \psi, t/\cos \psi)$.

- For the large volume stability conditions $Z_{(s,t)}^{LV}$, [Bousseau'19] constructed the scattering diagram \mathcal{D}_{ψ} in (s, t) upper half-plane for $\psi = 0$. For $\psi \neq 0$, just map $(s, t) \mapsto (s t \tan \psi, t/\cos \psi)$.
- The rays R(γ) are branches of hyperbola asymptoting to t = ±(s - d/r) for r ≠ 0, or vertical lines when r = 0. Walls of marginal stability W(γ, γ') are half-circles centered on real axis.



- For the large volume stability conditions $Z_{(s,t)}^{LV}$, [Bousseau'19] constructed the scattering diagram \mathcal{D}_{ψ} in (s, t) upper half-plane for $\psi = 0$. For $\psi \neq 0$, just map $(s, t) \mapsto (s t \tan \psi, t/\cos \psi)$.
- The rays R(γ) are branches of hyperbola asymptoting to t = ±(s - d/r) for r ≠ 0, or vertical lines when r = 0. Walls of marginal stability W(γ, γ') are half-circles centered on real axis.



• Think of $\mathcal{R}(\gamma)$ as the worldline of a fictitious particle of charge *r*, mass $m^2 = \frac{1}{2}d^2 - r \operatorname{ch}_2$ moving in a constant electric field !

Initial rays correspond to O(m) and O(m)[1], ie (anti)D4-branes with m units of flux, emanating from (s, t) = (m, 0) on the boundary where the central charge vanishes.



Initial rays correspond to O(m) and O(m)[1], ie (anti)D4-branes with m units of flux, emanating from (s, t) = (m, 0) on the boundary where the central charge vanishes.



 The first scatterings occur for t ≥ ¹/₂, after each constituent has moved by |Δs| ≥ ¹/₂. Causality and monotonicity of the 'electric potential' φ(γ) = d - sr along the flow, allow to bound the number and charges of constituents.



- {{ $-3\mathcal{O}(-2), 2\mathcal{O}(-1)$ }, \mathcal{O} }: 3 $\mathcal{O}(-2) \rightarrow 2\mathcal{O}(-1) \oplus \mathcal{O} \rightarrow E$ $K_3(2,3)K_{12}(1,1) \rightarrow -156$
- { $-\mathcal{O}(-3)$, { $-\mathcal{O}(-1)$, 2 \mathcal{O} }}: $\mathcal{O}(-3) \oplus \mathcal{O}(-1) \rightarrow 2\mathcal{O} \rightarrow E$ $K_3(1,2)K_{12}(1,1) \rightarrow -36$

Total:
$$\Omega_\infty(\gamma) = -192 = GV_4^{(0)}$$



- {{ $-\mathcal{O}(-5), \mathcal{O}(-4)$ }, $\mathcal{O}(-1)$ } $\mathcal{O}(-5) \rightarrow \mathcal{O}(-4) \oplus \mathcal{O}(-1) \rightarrow E$ $K_3(1,1)^2 \rightarrow 9$
- {{ $-\mathcal{O}(-4), \mathcal{O}(-3)$ }, { $-\mathcal{O}(-3), 2\mathcal{O}(-2)$ }} $\mathcal{O}(-4) \oplus \mathcal{O}(-3) \rightarrow$ $\mathcal{O}(-3) \oplus 2\mathcal{O}(-2) \rightarrow E$ $K_3(1,1)^2 K_3(1,2) \rightarrow 27$

•
$$\{-\mathcal{O}(-4), 2\mathcal{O}(-2)\}$$

 $\mathcal{O}(-4) \rightarrow 2\mathcal{O}(-2) \rightarrow E$
 $\mathcal{K}_6(1,2) \rightarrow 15$

Total: $\Omega_{\infty}(\gamma) = 51 = \chi(\text{Hilb}_4 \mathbb{P}^2)$

< ロ > < 同 > < 回 > < 回 >

- 2) Kähler moduli space of $K_{\mathbb{P}^2}$
- 3 Attractor flow trees and scattering diagrams
- 4 Large volume scattering diagram
- 5 Towards the exact scattering diagram

Exact scattering diagram

The full scattering diagram should interpolate between D^{LV}_ψ around τ = i∞ and D^o_ψ around τ = τ_o, and be invariant under the action of Γ₁(3).

Exact scattering diagram

- The full scattering diagram should interpolate between $\mathcal{D}_{\psi}^{\text{LV}}$ around $\tau = i\infty$ and \mathcal{D}_{ψ}^{o} around $\tau = \tau_{o}$, and be invariant under the action of $\Gamma_{1}(3)$.
- Under τ → τ/(3nτ+1) with n ∈ Z, O → O[n]. Hence we have an doubly infinite family of initial rays associated to O(m)[n].



Exact scattering diagram

- The full scattering diagram should interpolate between $\mathcal{D}_{\psi}^{\text{LV}}$ around $\tau = i\infty$ and \mathcal{D}_{ψ}^{o} around $\tau = \tau_{o}$, and be invariant under the action of $\Gamma_{1}(3)$.
- Under τ → τ/(3nτ+1) with n ∈ Z, O → O[n]. Hence we have an doubly infinite family of initial rays associated to O(m)[n].



• For $|\tan \psi| < \frac{1}{2\mathcal{V}}$ where $\mathcal{V} = \operatorname{Im} T(0) = \frac{27}{4\pi^2} \operatorname{Im} \operatorname{Li}_2(e^{2\pi i/3}) \simeq 0.463$ only the rays associated to $\mathcal{O}(m)[0]$ and $\mathcal{O}(m)[1]$ escape to $i\infty$, and merge onto rays in the large volume scattering diagram $\mathcal{D}_{\psi}^{\mathrm{LV}}$.

Exact scattering diagram - $\psi = 0$



B. Pioline (LPTHE, Paris)

BPS Dendroscopy

14/07/2022 30 / 35

< ロ > < 回 > < 回 >

In addition, there must be an infinite family of initial rays coming from τ = ^p/_q with q ≠ 0 mod 3, corresponding to Γ₁(3)-images of O(0).

- In addition, there must be an infinite family of initial rays coming from τ = ^p/_q with q ≠ 0 mod 3, corresponding to Γ₁(3)-images of O(0).
- This includes initial rays emitted at τ = n − ¹/₂, associated to Ω(n + 1); for ψ ~ ^π/₂, these merge onto initial rays of the orbifold scattering diagram.

- In addition, there must be an infinite family of initial rays coming from τ = ^p/_q with q ≠ 0 mod 3, corresponding to Γ₁(3)-images of O(0).
- This includes initial rays emitted at τ = n − ¹/₂, associated to Ω(n + 1); for ψ ~ ^π/₂, these merge onto initial rays of the orbifold scattering diagram.
- We conjecture that the only initial rays are the Γ₁(3) images of the structure sheaf O, each of them carrying Ω(kγ) = 1 for k = 1, 0 otherwise.

Exact scattering diagram - $\psi = \pm \frac{\pi}{2} \mod 2\pi$

• For $\psi = \pm \frac{\pi}{2}$, the diagram \mathcal{D}_{ψ}^{Π} simplifies dramatically, since the loci $\operatorname{Im} Z_{\tau}(\gamma) = 0$ are lines of constant $s := \frac{\operatorname{Im} T_D}{\operatorname{Im} T} = \frac{d}{r}$.



Exact scattering diagram - $\psi = \pm \frac{\pi}{2} \mod 2\pi$

• For $\psi = \pm \frac{\pi}{2}$, the diagram \mathcal{D}_{ψ}^{Π} simplifies dramatically, since the loci $\operatorname{Im} Z_{\tau}(\gamma) = 0$ are lines of constant $s := \frac{\operatorname{Im} T_D}{\operatorname{Im} T} = \frac{d}{r}$.



 Hence, there is no wall-crossing between τ_o and τ = i∞ when -1 ≤ d/r ≤ 0, explaining why the Gieseker index Ω_∞(γ) agrees with the index Ω_c(γ) in the anti-attractor chamber.

Exact scattering diagram, varying ψ

 $\gamma = [0, 1, 1) = \operatorname{ch} \mathcal{O}_{\mathcal{C}}:$



$\gamma = [1, 0, 1) = \operatorname{ch} \mathcal{O}$:



< 一型

-∢ ≣ ▶

 The scattering diagram is the proper mathematical framework for the attractor flow tree formula in the case of local CY3. This is because Z(γ) is holomorphic on M_K, so the gradient flow preserves the phase of Z(γ). Moreover, initial rays can only start from the boundary.

- The scattering diagram is the proper mathematical framework for the attractor flow tree formula in the case of local CY3. This is because Z(γ) is holomorphic on M_K, so the gradient flow preserves the phase of Z(γ). Moreover, initial rays can only start from the boundary.
- This provides an effective way of computing (unframed) BPS invariants in any chamber, and a natural decomposition into elementary constituents. Mathematically, different trees should correspond to different strata in *M_Z*(γ).

- The scattering diagram is the proper mathematical framework for the attractor flow tree formula in the case of local CY3. This is because Z(γ) is holomorphic on M_K, so the gradient flow preserves the phase of Z(γ). Moreover, initial rays can only start from the boundary.
- This provides an effective way of computing (unframed) BPS invariants in any chamber, and a natural decomposition into elementary constituents. Mathematically, different trees should correspond to different strata in M_Z(γ).
- It would be interesting to extend this description to other toric CY3, such as local del Pezzo surfaces, and to framed BPS indices.

< (□) ト < 三 ト

- The scattering diagram is the proper mathematical framework for the attractor flow tree formula in the case of local CY3. This is because Z(γ) is holomorphic on M_K, so the gradient flow preserves the phase of Z(γ). Moreover, initial rays can only start from the boundary.
- This provides an effective way of computing (unframed) BPS invariants in any chamber, and a natural decomposition into elementary constituents. Mathematically, different trees should correspond to different strata in M_Z(γ).
- It would be interesting to extend this description to other toric CY3, such as local del Pezzo surfaces, and to framed BPS indices.
- For a compact CY3, arg Z(γ) is no longer constant along the flow and there can be attractor points with Ω_{*}(γ) ≠ 0 at finite distance in Kähler moduli space...

< ロ > < 同 > < 回 > < 回 > < 回 > <

Thanks for your attention !



B. Pioline (LPTHE, Paris)

BPS Dendroscopy